

Trends in the Application of Dyadic Coping in Older Adults with Mild Cognitive Impairment and their Spouses

Yonghong Jiang¹; Lin Wang^{2*}; Lixia Peng¹

¹School of Nursing, Jinan University, China.

²Department of Transitional Care, The First Affiliated Hospital, Jinan University, China.

*Corresponding Author: Liling Wang

Email: wangliling@163.com

Abstract

Dyadic coping plays an important role in older adults with mild cognitive impairment and their spouses. Significant correlations were found between dyadic coping and self-efficacy, anxiety and depression, marital quality, and quality of life in older adults with mild cognitive impairment and their spouses. Psychosocial interventions, skills interventions, and exercise from the perspective of dyadic relationships were effective in improving the physical and mental health of older adults with mild cognitive impairment and their spouses. However, there is a lack of specific intervention programs for dyadic relationships in the local cultural context as an entry point. Therefore, it is necessary to draw on relevant internal and external literature to construct a dyadic coping intervention program suitable for older patients with mild cognitive impairment in China. The current article aims to provide a conceptual review focusing on dyadic coping care to inform the development of a dyadic intervention program suitable for older adults with mild cognitive impairment in China. This review outlines the theoretical concepts, assessment tools, current state of research, and intervention methods for mild cognitive impairment and dyadic coping.

Keywords: Mild cognitive impairment; Dyadic coping; Couples; Review.

Introduction

Mild Cognitive Impairment (MCI) is a clinical transition stage between cognitive normalcy and dementia Alzheimer's Disease (AD) with the main clinical manifestations of memory and executive function decline [1]. With the further aging process, up to 42.0% of the global elderly population over 60 years old is affected by mild cognitive impairment, and there are 9.83 million AD patients and 38.77 million MCI patients in the population over 60 years old in China [2]. Epidemiologic data published in the journal JAMA show that approximately 10% to 12% of patients with MCI progress to AD each year, with 80% developing AD after 6 years of follow-up [3]. Since the etiology and pathogenesis of AD are still unknown, there is no effective cure, and it will show irreversible degenerative changes, with

patients becoming progressively demented and incapacitated, which will bring great pain and economic burden to the family and the society; therefore, it is necessary to carry out early identification of and intervention in MCI at the pre-diagnostic stage of AD. Dyadic Coping (DC) refers to the joint responses and strategies of spouses in the face of stressful events [4], and studies have shown that the quality of interaction between patients and their spouses plays a crucial role in the treatment of chronic illnesses, and that their mental health and quality of life are interrelated [5]. Couples' supportive dyadic coping has a mediating effect in the relationship between patients' fear of disease progression, spousal anxiety, and marital quality, and the development of interventions using couples' supportive dy-

Citation: Jiang Y, Wang L, Peng L. Trends in the Application of Dyadic Coping in Older Adults with Mild Cognitive Impairment and their Spouses. Med Discoveries. 2024; 3(7): 1188.

adic coping as an entry point can help to improve the patients' and their spouses' negative emotions and improve marital quality [6]. Therefore, exploring dyadic coping between MCI patients and their spouses may provide a new perspective for early intervention management of MCI patients, which is important for slowing down the process of MCI and stopping its progression to dementia.

Conceptualization and assessment tools for mild cognitive impairment and dyadic coping

Concept of mild cognitive impairment and assessment tools

Mild cognitive impairment is a clinical condition in which an individual has an objectively measurable decline in memory or other cognitive domains that does not interfere with the ability to perform daily living tasks, and was first comprehensively described in 1997 by Petersen et al. [7]. Palmer et al. [8] proposed that MCI is a state of cognitive impairment between normal aging and dementia, and is usually categorized into amnesic MCI and non-MCI. Under the interaction of age, genetic factors, education level, lifestyle and other influencing factors, amnesic MCI is likely to progress to AD. The diagnosis of mild cognitive impairment in China often refers to the Chinese Expert Consensus 2021 on the Diagnosis and Treatment of Mild Cognitive Impairment of Alzheimer's Disease Origin [9], which includes: cognitive impairment reported by the patient or an informed person, or detected by an experienced clinician; the presence of objective evidence (neuropsychological tests) of impairment in 1 or more cognitive domains, with impairment in situational memory being the most common; complex instrumental everyday abilities can be mildly impaired but maintain independent daily living; diagnostic criteria for dementia have not been met. Diagnosis is often made with the help of neurological examination and neuropsychological assessments, such as the Montreal Cognitive Assessment (MoCA), the Activities of Daily Living Scale (ADL), the Hamilton Hamilton Anxiety Scale/Hamilton Depression Scale, etc.

The concept of dyadic coping and assessment tools

German scholar Bodenmann [10] first proposed the concept of dyadic coping in the early 1990s, referring to the ability of partners to jointly cope with problem solving in the face of dyadic stressful events. Mariana et al. [11] proposed that the dyadic coping process consists of: clear communication of stress; individual positive/negative coping; joint positive/negative coping, after integrating the process of dyadic coping. Randomized controlled studies of dyadic coping were conducted earlier abroad, and scholars have gradually shifted the study of the disease from an individualized to a dyadic perspective. A study investigated 154 pairs of men with prostate cancer and their spouses, and found that when dealing with cancer-related stress, the support and understanding between husband and wife helped to alleviate the patient's psychological distress. Sterba et al. [12] found that higher dyadic efficacy helped to increase the success rate of smoking cessation among smokers. Domestic nursing research on dyadic coping started late, Luo Qun [13] took the lead in exploring the association between dyadic coping styles and intimacy between gynecologic cancer patients and their spouses, and the results showed that the dyadic supportive coping of gynecologic cancer patients and spouses was related to each other, and the intimacy of the spouses and patients was related to each other; Conceivably, interventions centered on the coping styles of couples are an important way for couples to strengthen each other in the face of the challenges of cancer.

An important approach to strengthen couples' resources in the face of cancer challenges. In recent years, scholars at home and abroad have tried to use the dyadic coping theory in the management of patients and spouses with cancer, chronic diseases, and obstetrics and gynecology [5,14,15].

There are no standardized criteria for assessing dyadic coping, and the most widely used assessment tool is the Dyadic Coping Inventory (DCI) developed by Gmelch et al. [16] in 2008. The scale consists of six dimensions, namely stress communication, positive coping, negative coping, empowering coping, coping together, and quality evaluation, with a total of 37 entries. The scale is scored from 1 to 5, representing "rarely" to "very frequently", and the higher the score, the higher the level of dyadic coping, which was Chineseized by our scholars Xu et al. [17] in 2016. In addition, the Revised Dyadic Adjustment Scale (RDAS) and the Locke-Wollance Marital Adjustment Scale (LWMS) are also commonly used dyadic coping assessment tools [18,19].

Current status of research on dualistic coping in older adults with MCI and their spouses

Self-efficacy refers to an individual's ability and beliefs when faced with a wide range of task demands or completely new situations [20]. Tonga et al. [21] found in a cross-sectional study that positive mood could be enhanced by increasing patients' confidence in their ability to cope with their illness and daily life, thereby improving overall quality of life. As for spouses of patients, increasing their self-efficacy not only helps them to better cope with the caregiving burden of AD or other types of cognitively impaired disorders, but also reduces depressive symptoms [22], significantly improves caregiving outcomes, and slows down the decline of cognitive functioning levels in the elderly [23].

The correlation between dyadic coping and self-efficacy in older adults with MCI

The Chinese version of the General Self-Efficacy Scale (GSES) was co-developed by Zhang and Schwarze [24], which contains 10 entries on a four-point scale, with higher mean scores indicating higher self-efficacy for individuals. Kurasz et al. [25] adapted a self-efficacy scale used for the management of chronic illnesses to make it applicable to patients with Mild Cognitive Impairment (MCI) and more targeted. Furthermore, he suggested that self-efficacy, especially memory self-efficacy (memory-SE), may be a key factor in maintaining patients' cognitive function and psychosocial integrity. Domestic scholars Tian Si-Ying et al. [26] found that MCI patients' self-efficacy was reduced after memory loss, and they were prone to self-doubt, self-denial and self-complaints, lack of confidence in many things, and loss of goals and motivation in life. In recent years, research on the relationship between dyadic coping and self-efficacy has achieved certain results, and the mediating roles played by the two in different diseases have been gradually clarified. Wang et al. [27] selected 206 spouses of patients with cervical cancer from five hospitals in Jiangsu Province, and used the 12-item Brief Health Questionnaire (SF-12), the General Self-Efficacy Scale, and the Dyadic Coping Inventory (DCI) were investigated and structural equation modeling was constructed. The results showed that self-efficacy partially mediated between dyadic coping and quality of life, and the mediating effect accounted for 16.0% of the total effect. Hui-Ying An [6] used purposive sampling to conduct semi-structured interviews with 12 pairs of young and middle-aged lymphoma patients and their spouses to gain in-

sights into the couples' experience of disease coping and to explore the positive factors that promote the couples' adaptation to the disease. The results showed that patients' and spouses' dyadic coping status had different degrees of mediating effects in the relationship between their self-efficacy, negative emotions, and marital quality.

Therefore, patients and spouses should be regarded as a coping whole, focusing on dyadic strategies such as stress communication and supportive coping, and carrying out couple-centered coping interventions from the perspective of improving self-efficacy and negative emotions of both partners, so as to improve the quality of marriage, and then improve the quality of life of both partners. However, no research has been conducted on the mediating effects of dyadic coping, self-efficacy, quality of life, and negative emotions on MCI patients and their spouses, and it is necessary to construct a structural equation model to further clarify the relationship between them, so as to provide practical guidance for mental health research and clinical work, and to provide reference for preventing and delaying the transformation of dementia.

The correlation between dyadic coping and anxiety and depression in older adults with MCI

Most previous studies have focused only on the mental health status of patients with MCI, and fewer studies have analyzed the emotional problems of their spouses. As early as 2005, in 1 cross-sectional study, researchers Garand et al. [28] found that even in the early stages of the onset of MCI, spouses of patients with MCI already face a caregiver burden similar to that of dementia caregivers, and there is an incremental mental health risk that progresses over time. In 2017, the World Health Organization established caregiver needs as one of the 2017-2025 one of the priorities of the Dementia Patient Care Plan [29]. Scholars Kim et al. [30] selected 2,782 men and 2,515 women, totaling 5,307 study participants from the Korean Longitudinal Study of Aging (K-LSA) database, and assessed the depressive symptoms of the study participants with the Center for Epidemiologic Studies Depression Scale (CES-D 10), and with the Korean version of the Brief Mental State Examination Scale (K-MMSE) to evaluate the cognitive function of their spouses. The results showed that study subjects living with spouses with mild dementia (MMSE 18-23)/moderate to severe dementia (MMSE \leq 17) exhibited higher depressive symptoms (CES-D \geq 4) compared to those living with spouses with normal cognitive functioning; furthermore, those who did not engage in social activities had higher depression scores than those who did.

The role of spouses in the maintenance of cognitive function in older adults with MCI

In a cross-sectional study of the relationship between dyadic coping and Quality of Life (QoL) in patients with dementia and their spouses, Schumann et al. [31] found that patients had higher QoL scores when their spouses were the primary caregivers compared with when their children were the primary caregivers. Yan Pingting et al. [32] conducted a cross-sectional survey of 8,221 residents aged 65 years and older in urban and rural areas of Hubei Province, using face-to-face interviews to complete general surveys and neuropsychological assessments, as well as clinical physical examinations and laboratory tests, and diagnostic expert panels to diagnose MCI and dementia in participants. The results showed that the prevalence of MCI and dementia increased in the nulliparous elderly group, especially for men aged 65-69 years and 80 years and older, and that

having a spouse is essential to protect cognitive function in the elderly. The significant association between marital status and the risk of dementia and mild cognitive impairment is consistent with the findings of Chen Lu, Liu et al., Su Yuan, Skirbekk et al. [33-36]. Zhao Xiaohang et al. [37] also suggested that widowhood affects cognitive functioning and mental health in older adults; in terms of basic daily living skills, for men with high marital quality, their Instrumental Activities of Daily Living (IADL) skills such as housework, cooking, purchasing, and handling personal For men with high marital quality, their IADL such as housework, cooking, purchasing, and handling personal things decreased significantly, while for women, marital quality had no significant effect on their IADL abilities after widowhood. Scholars Yu (Yu Yifan) et al. [38] examined the relationship between living arrangements and cognitive decline in Chinese older adults, and further examined gender differences on these relationships. It was found that cognitive decline was slower in older adults who lived with their spouses, and that women had a smaller effect on their cognitive decline when living alone compared to men. This suggests that spouses contribute to the improvement of cognitive decline in people with MCI, and more so for men, and that appropriate responses can be developed by considering the couple as a whole and advancing public policies to support caregiving families.

Current status of interventions for dyadic coping in older adults with MCI and their spouses

Dyadic psychosocial interventions

Many treatment programs are often carried out for individual patients or caregivers, but a meta-analysis showed that interventions are less effective when they are targeted only to the individual [39]. A systematic evaluation by Poon et al. [40] showed that Cognitive Stimulation Therapy (CST) is currently mostly applied in studies of psychological interventions for people with MCI. This therapy was first pointed out by the clinical guidelines issued by the National Institute for Clinical Excellence in the UK, which can be used as a non-pharmacological intervention to treat dementia, emphasizing the suppression and abatement of the various aspects and stages of wrongly acquired behaviors, and the correction of problematic behaviors for the treatment of mental illnesses [41]. Yates et al. [42], in a Randomized Controlled Trial (RCT), regarded patients with dementia and their family caregivers as a whole, and made a CST modified to form Individual Cognitive Stimulation Therapy (iCST), which was family caregiver-led and dementia patient-centered, and 356 intervened with the subjects. Results showed that although people with dementia treated with iCST did not significantly improve cognitive function/quality of life, and iCST was not found to improve caregivers' mental or physical health, the therapy had a positive impact on improving caregiving relationships and improving caregivers' quality of life, suggesting that iCST may be an important component of in-home care for people with dementia in order to help them receive effective long-term care. This was also confirmed by Myhre et al. [43] study on the coping experience of spouses of patients with dementia, where spouses who participated in Norwegian Cognitive Behavioral Therapy for Early Stage Dementia (N-CORDIAL) with their dementia patients showed an improvement in coping strategies 6-12 months post-intervention and were able to utilize memory aids such as calendars/memos, as well as a positive emotional orientation to cope with the patient.

Spouses, as the primary caregivers of patients with MCI/de-

mentia, play a key role in the provision of in-home care; therefore, strengthening family resources is an important task for caregivers, and further research is needed on how to effectively educate caregivers about the disease, and develop training in problem solving and dealing with negative emotions in order to improve the coping strategies of spouses of patients with MCI/dementia in the long term.

Dyadic skills interventions

Greenaway et al. [44] conducted a Memory Support System (MSS) cognitive rehabilitation intervention with 40 patients with MCI, in which patients compensated for their memory loss by using calendars/notebooks, etc., to record to-do lists and journaling about tasks that needed to be accomplished at a specific time or day. Participants in the intervention group received a total of 12 one-hour sessions of MSS training for 6 weeks. The training consisted of instruction, demonstration, practice, and assignment of homework. The control group was only provided with the MSS calendar/notebook and encouraged to use it on their own, but had no further verbal or written instruction. Results showed that MCI patients who received MSS training showed significant improvements in Activities of Daily Living (ADLs) and memory self-efficacy, and improvements in ADLs in the intervention group compared with the control group who were only provided with calendars/notebooks were sustained until the follow-up visit 8 weeks later. In addition, spouses in the intervention group had significantly lower Center for Epidemiological Survey, Depression scale (CES-D) scores than spouses in the control group at the 8-week and 6-month post-intervention follow-ups. This suggests that MSS training helps patients to better manage their daily lives and reduces the frequency with which spouses need to intervene, thus improving their mood and quality of life. Subsequently, a study by Khayoun et al. [45] also demonstrated that positive emotional feedback from spouses, as well as support and assistance provided to patients with MCI, can help patients to successfully adapt to the stress of the disease, and their results showed that partner involvement and support were crucial to the patients' learning outcomes of the MSS, and that the higher the partner's openness, the greater the willingness of the patients to learn the MSS, and the shorter the time it took to master the MSS.

In a randomized controlled trial by researcher Lu et al. [46], 36 pairs of MCI patients and their spouses were intervened, of which 17 pairs received the Daily Enhancement of Meaningful Activity (DEMA) intervention and 19 pairs received the Attention Control Intervention. The DEMA program is a skills training program for MCI patients and their spouses, and their spouses, a skills-training program designed to help couples learn how to supportive dyadic coping to adapt to possible changes in the condition as it progresses over time. It consists of 6 offline sessions of 1 hour each, conducted once every 2 weeks, covering 6 self-management needs: planning meaningful activities; understanding MCI; treatments for MCI; understanding and coping with negative emotions; learning to live with and cope with MCI; how to seek out healthcare resources that can give help. The results showed that the DEMA group had higher bidirectional consistency in perceived functional ability and life satisfaction at 3 months post-intervention and improved somatic functioning at 2 weeks post-intervention compared to the attention control intervention control group.

Dyadic movement interventions

Ahn et al. [47] conducted a 16-week intervention with four

pairs of patients with Mild Cognitive Impairment (MCI) and their spouses to assess the role of couples engaging in physical activity together in improving physical health and psychological well-being of both partners. The study included aerobic exercise and muscle strength training, and the results showed that patients with MCI had improved partner support scores after the intervention, exercising with their partners enhanced mutual support and encouragement; aerobic fitness was improved by an increase in the 6-minute walk test distance; fall efficacy scale scores were reduced, and the fear of falling was alleviated; the time to complete the color-vocabulary interference task was shortened, executive functioning improved; quality of life scores improved. Spouses' positive affect scores increased, Zarit Caregiver Burden Scale scores decreased, and psychological well-being increased.

Conclusion

There is a correlation between the dyadic coping strategies employed by patients with mild MCI and their spouses and the indicators of physical and mental health outcomes. These include improvements in self-efficacy, reductions in depression and anxiety symptoms, and delays in cognitive decline, among others. Intervention studies utilising dyadic relationships as a starting point can effectively enhance the level of psychological adaptation, subsequently enhancing the quality of life. In China, research on dyadic coping commenced relatively late, with the majority of cross-sectional studies conducted. Furthermore, interventions for patients with MCI have been limited to the individual patient, with couples not being considered as a whole in the treatment programme. In future research, we need to drawing on mature foreign experience and combining it with our culture, conduct high-quality randomised controlled trials, track the effects of the interventions on a regular basis, and develop dyadic coping interventions adapted to China's MCI geriatric population, so as to improve the mental health outcomes of both spouses and thus enhance their quality of life. In addition, we need to actively develop "Internet + Nursing", refers to medical institutions using Internet technologies and big data to provide nursing services to discharged patients or those with severe illnesses unable to visit hospitals, through online applications and offline care provision. "Internet + Nursing" can break through the previous limited verbal health education, conduct home skills training, and strengthen WeChat/telephone/door-to-door follow-up visits, so as to provide MCI couples with appropriate health education in a timely, efficient and convenient manner, thus improving their ability to cope with the disease and slowing down the disease process. Pay attention to rural areas with relatively insufficient medical and health resources and a relatively low level of education. This can be achieved by increasing the sample size and expanding the scope of the survey, which will provide a theoretical basis for relevant epidemiological studies. At the same time, it is necessary to guide high-quality medical resources down to the grassroots level, intensify screening and intervention efforts, and provide more comprehensive psychosocial support and treatment programmes for MCI patients and their spouses.

Funding: Guangdong Medical Science and Technology Research Fund Project, No. A2022389.

References

1. L Jia, Y Du, L Chu, Z Zhang, F Li, et al. Prevalence, risk factors, and management of dementia and mild cognitive impairment in adults aged 60 years or older in China: a cross-sectional study.

- The Lancet public health. 2020; 5: e661-e671.
2. R Ren, J Qi, S Lin, X Liu, P Yin, et al. The China alzheimer report 2022. *General psychiatry*. 2022; 35.
 3. Langa K M, Levine D A. The diagnosis and management of mild cognitive impairment: a clinical review. *Jama*. 2014; 312: 2551-2561.
 4. BAI Yang, FENG Xianling, GU Yujun, LI Xiaoliu. Study on the Relationship Among Dyadic Coping, Marital Adjustment and Quality o-f Life in Patients with Cervical Cancer Undergoing Chemotherapy Base-d on the Actor-Partner Interdependence Model. *Military Nursing*. 2024; 41: 65-69.
 5. Vagharseyyedin SA, Arbabi M, Rahimi H, Moghaddam SG. Effects of a caregiver educational program on interactions between family caregivers and patients with advanced COPD. *Home Healthcare Now*. 2022; 40: 75-81.
 6. An Huiying, Lei Lei, Cui Panpan. Relationship among negative emotions, marital quality, and supportive duality coping in Relationship among negative emotions, marital quality, and supportive duality coping in young and middle-aged patients with lymphoma and their spouses. *Chinese Mental Health Journal*. 2023; 37: 1059-1064.
 7. Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, et al. Mild cognitive impairment: clinical characterization and outcome. *Archives of neurology*. 1999; 56: 303-8.
 8. Palmer K, Fratiglioni L, Winblad B. What is mild cognitive impairment? Variations in definitions and evolution of nondemented persons with cognitive impairment. *Acta Neurologica Scandinavica*. 2003; 107: 14-20.
 9. Dementia and Cognitive Disorders Group of the Neurology Section of the Chinese Medical Association. Chinese expert consensus on the diagnosis and treatment of mild cognitive impairment of Alzheimer's disease origin 2021. *Chinese Journal of Neurology*. 2022; 55: 20.
 10. Bodenmann G. Dyadic coping: a systemic-transactional view of stress and coping among couples: Theory and empirical findings. *Revue Européenne De Psychologie Appliquée*. 1997; 47: 137-141.
 11. Bantia R, Malcarne VL, Varni JW, Ko CM, Sadler GR, et al. The effects of dyadic strength and coping styles on psychological distress in couples faced with prostate cancer. *Journal of behavioral medicine*. 2003; 26: 31-52.
 12. Sterba KR, Rabius V, Carpenter MJ, Villars P, Wiatrek D, et al. Dyadic efficacy for smoking cessation: preliminary assessment of a new instrument. *Nicotine & Tobacco Research*. 2011; 13: 194-201.
 13. Luo Qun. The relationship between dyadic coping and intimacy of patient switch gynecological cancer and their spouses. *Anhui Medical University*. 2017.
 14. von Heymann-Horan A, Bidstrup PE, Johansen C, Rottmann N, Andersen EA, et al. Dyadic coping in specialized palliative care intervention for patients with advanced cancer and their caregivers: Effects and mediation in a randomized controlled trial. *Psycho Oncology*. 2019; 28: 264-70
 15. Wang Jing, Zhang Haiyang, XU Yanyu, Yu Shengsheng, Yan Ying, et al. Effects of the dyadic coping nursing intervention for treating patients with ovarian cancer after operation on their self-man-agement ability and post-traumatic growth. *Chinese Journal of Family Planning*. 2024; 32: 589-594.
 16. Gmelch S, Bodenmann G, Meuwly N, Ledermann T, Steffen Sozinova O, et al. Dyadisches Coping Inventar (DCI): Ein Fragebogen zur Erfassung des partnerschaftlichen Umgangs mit Stress. *Zeitschrift für Familienforschung*. 2008; 20: 185-203.
 17. Xu F, Hilpert P, Randall AK, Li Q, Bodenmann G. Validation of the Dyadic Coping Inventory with Chinese Couples: Factorial Structure, Measurement Invariance, and Construct Validity. *Psychological Assessment*. 2016; 28: e127.
 18. Antoine P, Christophe V, Nandrino JL. Dyadic Adjustment Scale: Clinical interest of a revision and validation of an abbreviated form. *L Encéphale*. 2008; 34: 38-46.
 19. Wang Q, Wang D, Li C, Miller RB. Marital satisfaction and depressive symptoms among Chinese older couples. *Aging Ment Health*. 2014; 18: 11-18.
 20. Schwarzer R. Optimistic self-beliefs: Assessment of general perceived self-efficacy in thirteen cultures. *World Psychology*. 1997; 3: 177-190.
 21. Tonga JB, Eilertsen DE, Solem IK, Arnevik EA, Korsnes MS, et al. Effect of self-efficacy on quality of life in people with mild cognitive impairment and mild dementia: themediating roles of depression and anxiety. *American Journal of Alzheimer's Disease & Other Dementias®*. 2020; 35: 1533317519885264.
 22. Langer K, O'Shea DM, De Wit L, DeFeis B, Mejia A, et al. Self-efficacy mediates the association between physical function and perceived quality of life in individuals with mild cognitive impairment. *Journal of Alzheimer's Disease*. 2019; 68: 1511-1519.
 23. Khan TS, Hirschman KB, McHugh MD, Naylor MD. Self-efficacy of family caregivers of older adults with cognitive impairment: a concept analysis. *Nurs Forum*. 2021; 56: 112-126.
 24. Zhang J X, Schwarzer R. Measuring optimistic self-beliefs: a Chinese adaptation of the general self-efficacy scale. *Psychologia: An International Journal of Psychology in the Orient*. 1995.
 25. Kurasz AM, DeFeis B, Locke DE, De Wit L, Amofa P, et al. Psychometric properties of the self-efficacy for managing mild cognitive impairment scale. *International Journal of Geriatric Psychiatry*. 2021; 36: 174-181.
 26. TIAN Siying, CHANG Hong, LI Peipei, ZHANG Xinyue, QIAO Yuchen, et al. A qualitative study on emotional experience of patients with mild cognitive impairment. *Chinese Journal of Nursing*. 2022; 57: 1206-1212.
 27. Wang C, Wacharasin C, Hengudomsub P. Self-efficacy as mediators of the association between dyadic coping and quality of life among spousal caregiver of patients with cervical cancer. *Asia-Pacific Journal of Oncology Nursing*. 2024; 11: 100354.
 28. Garand L, Amanda Dew M, Eazor LR, DeKosky ST, Reynolds III CF. Caregiving burden and psychiatric morbidity in spouses of persons with mild cognitive impairment. *International journal of geriatric psychiatry*. 2005; 20: 512-522.
 29. World Health Organization. Global action plan on the public health response to dementia 2017–2025. *World Health Organization*. 2017.
 30. Kim W, Lee TH, Shin J, Park EC. Depressive symptoms in spouse caregivers of dementia patients: A longitudinal study in South Korea. *Geriatrics & Gerontology International*. 2017; 17: 973-983.
 31. Schumann C, Alexopoulos P, Perneckzy R. Determinants of self- and carer-rated quality of life and caregiver burden in Alzheimer disease. *Int J Geriatr Psychiatry*. 2019; 34: 1378-1385.

32. Yan Pingting, Liu Dan, Cheng Guirong, Shi Chongqing, Huang Jing, et al. Association of Late-life Marital Status with Mild Cognitive Impairment and Dementia in Old Age. *Journal of Sun Yat-sen University (Medical Sciences)*. 2022; 43: 229-237.
33. Chen Lu. Study on the Impacts of Marital Status on the Health of the Elderly in China. Zhejiang University. 2016.
34. Liu H, Zhang Y, Burgard SA, Needham BL. Marital status and cognitive impairment in the United States: evidence from the National Health and Aging Trends Study. *Annals of Epidemiology*. 2019; 38: 28-34.
35. Su Yuan. A Study of the Impact of Marital Status on the Health of the Elderly in the Context of Active Ageing-Based on the Analysis of CLHLS Data. Guizhou University. 2022.
36. Skirbekk V, Bowen CE, Håberg A, Jugessur A, Engdahl B, et al. Marital histories and associations with later-life dementia and mild cognitive impairment risk in the Hunt4 70+ study in Norway. *Journal of aging and health*. 2023; 35: 543-55.
37. Zhao Xiaohang, Li Jianxin. Impact of Widowhood on the Health of Chinese Older Adults: The Moderating Role of Social Connectedness. *Population Journal*. 2022; 1.
38. Yu Y, Lv J, Liu J, Chen Y, Chen K, et al. Association between living arrangements and cognitive decline in older adults: a nationally representative longitudinal study in China. *BMC geriatrics*. 2022; 22: 843.
39. Orrell M, Bebbington P. Life events and senile dementia. I. Admission, deterioration and social environment change. *Psychological medicine*. 1995; 25: 373-386.
40. Poon E. A systematic review and meta-analysis of dyadic psychological interventions for BPSD, quality of life and/or caregiver burden in dementia or MC. *Clinical gerontologist*. 2022; 45: 777-797.
41. National Institute for Clinical Excellence. Social Care Institute for Excellence (NICE/SCIE). *Dementia: supporting people with dementia and their careers in health and social care*. London: National Institute for Health and Care Excellence. 2006.
42. Yates LA, Leung P, Orgeta V, Spector A, Orrell M. The development of individual cognitive stimulation therapy (iCST) for dementia. *Clinical interventions in aging*. 2015; 30: 95-104.
43. Myhre J, Bjørnstad Tonga J, Ulstein ID, Høye S, Kvaal K. The coping experiences of spouses of persons with dementia. *Journal of clinical nursing*. 2018; 27: e495-e502.
44. Greenaway M C, Duncan N L, Smith G E. The memory support system for mild cognitive impairment: randomized trial of a cognitive rehabilitation intervention. *International journal of geriatric psychiatry*. 2013; 28: 402-409.
45. Khayoun R, Devick KL, Chandler MJ, Shandera-Ochsner AL, De Wit L, et al. The impact of patient and partner personality traits on learning success for a cognitive rehabilitation intervention for patients with MCI. *Neuropsychological Rehabilitation*. 2022; 32: 2483-95.
46. Lu YY, Bakas T, Yang Z, Weaver MT, Austrom MG, et al. Feasibility and effect sizes of the revised daily engagement of meaningful activities intervention for individuals with mild cognitive impairment and their caregivers. *Journal of gerontological nursing*. 2016; 42: 45-58.
47. Ahn S, Cobb SJ, Crouter SE, Lee CE, Crane MK, et al. Physical activity together for couples living with mild cognitive impairment (PAT-MCI): A feasibility study. *Geriatric Nursing*. 2024; 55: 221-8.